

Examples of splints for the management of carpal tunnel syndrome



For further information and assistance contact your:

- General Practitioner
- Woman's Health Physiotherapist

King Edward Memorial Hospital
Physiotherapy Department (Subiaco)

Telephone: (08) 9340 2790

Disclaimer: The advice and information contained herein is provided in good faith as a public service. However the accuracy of any statements made is not guaranteed and it is the responsibility of readers to make their own enquiries as to the accuracy, currency and appropriateness of any information or advice provided. Liability for any act or omission occurring in reliance on this document or for any loss, damage or injury occurring as a consequence of such act or omission is expressly disclaimed.

Compiled by: Physiotherapy Department

Email: ogccu@health.wa.gov.au

Website: <http://wnhs.health.wa.gov.au>

Produced by: Women and Newborn Health Service

© November 2008 WNHS 0513

Reviewed October 2010

This information is available in alternative formats upon request

WOMEN AND NEWBORN HEALTH SERVICE

King Edward Memorial Hospital

374 Bagot Road Subiaco WA 6008

Telephone: (08) 9340 2222



Government of Western Australia
Department of Health



Delivering a Healthy WA

Carpal Tunnel Syndrome in Pregnancy



What is Carpal Tunnel Syndrome (CTS)?

Carpal tunnel syndrome is a disorder of the hand caused by pressure on the median nerve as it runs through the wrist.

Carpal tunnel syndrome may occur due to overuse and repetitive hand movements, arthritis or during pregnancy. In pregnancy hormonal changes may result in fluid retention, which can compress the median nerve. CTS triggered during pregnancy usually resolves soon after birth.

CTS most frequently occurs during the third trimester, but it may occur during the first and second trimesters.

What is the cause of CTS in pregnancy?

Swelling in the hand/ wrist caused by fluid retention compresses the median nerve.

Symptoms may be exacerbated by repetitive hand movements, holding hands in sustained positions or by putting weight through outstretched hands, as in kneeling on all fours.

What are the signs and symptoms of CTS?

- Numbness
- Pins and needles especially in the thumb, index and middle fingers
- Symptoms may worsen or only occur at night
- Radiating or referred pain into the forearm or upper arm
- Weakness of the hand/ weak grasp
- Pain and problems with thumb movements
- Clumsiness and loss of hand coordination




Figure 1. Tendon gliding exercises

What are the treatment options for CTS?

- Rest/ Modify activities
- Massage
- Contrast baths (alternating hands in warm and cold water)
- Ice
- Elevation
- Splints (working and resting, especially at night)
- Muscle pump exercises/ gliding exercises
- Avoid sleeping on the affected side
- Electrotherapy - ultrasound
- Cortisone injection into the wrist
- Surgery as a last resort




Figure 2. Nerve gliding exercises