

CLINICAL PRACTICE GUIDELINE

Guideline coverage includes NICU KEMH, NICU PMH and NETS WA

Transillumination of the Chest

This document should be read in conjunction with the [Disclaimer](#)

To detect abnormal air collections within the chest cavity in an infant suspected of having an air leak or any respiratory supported infant who deteriorates with asymmetrical chest wall movement or air entry.

Transillumination can be done by any staff member deemed competent or a trainee under direct supervision from a competent staff member. If an abnormal air collection within the chest cavity is suspected, a Medical Officer competent in needle aspiration of the chest must be contacted immediately to attend the unit.

Any suspicion of an accumulation of air by transillumination should be confirmed by a chest X-ray if time permits.

[Emergency needle aspiration](#) equipment is kept in a container at each infant's bay for NICU infants.

For infants suspected of having fluid in the chest cavity, refer to [ICC insertion](#).

Equipment

- High intensity fiber-optic cold light source (transilluminator).
- Cardiopulmonary monitoring.
- Blood pressure monitoring.

Procedure

1. Lower the lights in the room to enable hyper-lucent areas to be seen if present.
2. Place the transilluminator along the posterior axillary line on the side on which the air collection is suspected. The transilluminator may be moved up and down along the posterior axillary line and above the nipple to detect any areas of increased transmission of light.
3. Place the transilluminator in the third or fourth intercostal space on the left midclavicular line and angle the light towards the xiphoid process to detect any areas of increased transmission of light.
4. Transilluminate both sides of the chest to give a comparison.



Image: Transillumination of the chest

Note:



- A **false-positive** result may occur with severe subcutaneous chest wall oedema and pulmonary interstitial emphysema.
- A **false-negative** result may occur in infants with a thick layer of subcutaneous fat.

References

1. Karlsen, K. Post-resuscitation / pre-transport stabilization care of sick infants: guidelines for neonatal healthcare providers. 6th Edition. Park City, Utah. 2013.

Related WNHS policies, procedures and guidelines

Neonatal Clinical Guidelines: - [Needle aspiration of the Chest](#)
- [Intercostal Catheter Insertion](#)

| | | | |
|-----------------------|---|-------------------|-----------------------------|
| Document owner: | Neonatal Directorate Management Committee | | |
| Author / Reviewer: | Neonatal Directorate Management Committee | | |
| Date first issued: | June 2006 | | |
| Last reviewed: | 11 th April 2017 | Next review date: | 11 th April 2020 |
| Endorsed by: | Neonatal Directorate Management Committee | Date endorsed: | 27 th June 2017 |
| Standards Applicable: | NSQHS Standards: 1  Governance, 9  Clinical Deterioration | | |

**Printed or personally saved electronic copies of this document are considered uncontrolled.
Access the current version from the WNHS website.**