



Clinical Practice Guideline

Guideline coverage includes NICU KEMH, NICU PMH and NETS WA

Resuscitation: Who Attends Births (KEMH)

This document should be read in conjunction with the [Disclaimer](#)

The decision to call personnel for possible resuscitation of the newborn is the responsibility of the obstetric staff. Sufficient notice should be given to enable staff members from the Department of Neonatal Paediatrics to get to the delivery ward or theatre, to check and prepare the resuscitation trolley and to obtain a resume of the relevant pregnancy and labour details.

A neonatal consultant will attend high risk deliveries if requested by the neonatal SR or obstetric consultant. A routine caesarean section is not to commence until the relevant NICU staff are present in theatre.

These lists are not exclusive. If in doubt call for help.

<p>GROUP ONE RMO's - Neonatal registrars will support RMO's at every birth until the RMO is deemed competent and confident in attending births alone.</p>	<p>GROUP TWO RMO's and Neonatal Registrar - a SR/Consultant should be called to attend at the birth if needed.</p>
<ul style="list-style-type: none"> • < 37 weeks gestation • Forceps (low cavity) • Vacuum extraction (low cavity) • Pre-eclampsia • Intrauterine growth retardation • Membranes ruptured greater than 24hours • Maternal sepsis/positive maternal HVS for GBS • Maternal diabetes • All elective/non complicated caesarean sections at term and under regional anaesthesia • Maternal morphine analgesia given within 4 hours of birth 	<ul style="list-style-type: none"> • < 35 weeks gestation • All vaginal breech births and multiple pregnancies • Poor obstetric history • High or mid cavity forceps / vacuum extraction • Trial of instrumental birth in theatre • Meconium liquor, foetal bradycardia and other foetal distress • Rhesus iso-immunisation • APH/Intrapartum bleed • Caesarean section that is non-elective or under general anaesthetic


GROUP THREE (HIGH RISK): RMO's and Neonatal Registrar and notify SR and/or Neonatal Consultant.

A resuscitation cot with full intensive care facilities is needed.

- Code 'Blue' paediatric
- All deliveries < 30 weeks gestation and other high risk births
- Multiple pregnancy < 34 weeks gestation
- Severe foetal distress
- Severe rhesus iso-immunisation e.g. Hydrops
- Known high risk congenital anomalies such as diaphragmatic hernia

Related WNHS policies, procedures and guidelines

Neonatology Clinical Guidelines – [Resuscitation Algorithm for the Newborn](#)
[Resuscitation: Neonatal](#)
[Resuscitation: Medications and Fluids](#)

Document owner:	Neonatology Directorate Management Committee		
Author / Reviewer:	Neonatology Directorate Management Committee		
Date first issued:	October 2013		
Last reviewed:	27 th September 2016	Next review date:	27 th September 2016
Endorsed by:	Neonatology Directorate Management Committee	Date endorsed:	27 th September 2016
Standards Applicable:	NSQHS Standards: 1  Governance, 6  Clinical Handover, 9  Clinical Deterioration		

**Printed or personally saved electronic copies of this document are considered uncontrolled.
 Access the current version from the WNHS website.**