



Government of **Western Australia**
Department of **Health**

WOMEN AND NEWBORN HEALTH SERVICE

King Edward Memorial Hospital

G.P. INFORMATION AND FOLLOW-UP LETTER

Date:...../...../.....

ADDRESSOGRAPH

Dear Doctor.....,

Thank you for reviewing.....and family. This neonate was identified as at risk for perinatal transmission of Hepatitis B virus (HBV).

The mother of this infant was diagnosed as Hepatitis B virus positive. The neonate has received:

- 1st dose of hepatitis B vaccine
- Hepatitis B immunoglobulin

It is recommended the infant receives completion of the vaccination course for prevention of HBV recommended according to the Australian Immunisation Handbook for 2008.

Recommended follow-up testing for the infant at risk of perinatal infection for HBV is as follows:

- Perform bloods for Anti-HBs and HBsAg levels 3-12 months following the primary vaccination course.
- If anti-HBs levels are adequate and the HBsAg is negative, then the child is considered immune through natural clearance or vaccination.

If this schedule is followed more than 90% of babies born to mothers who are HBV positive will not develop infection.

Breastfeeding is not contra-indicated for neonates born to mothers who are hepatitis B surface antigen (HBsAg) positive.

Thank you for your ongoing care of this family,

Regards

PRINT NAME:..... Signature:..... Designation:.....

Staff of NCCU, King Edward Memorial Hospital