



CLINICAL PRACTICE GUIDELINE

Hyperemesis Gravidarum

This document should be read in conjunction with the [Disclaimer](#)

Aim

The appropriate referral and management of hyperemesis for women booked into the Community Midwifery Program.

For background Information consult the KEMH Clinical Guideline [Hyperemesis Gravidarum](#)

- Inform the woman of the need for consultation and referral to the support hospital and/or G.P for management.

References





1. American College of Obstetrics and Gynecology, authors. ACOG (American College of Obstetrics and Gynecology) Practice Bulletin: nausea and vomiting of pregnancy. **Obstet Gynecol.** 2004;103:803–814.
2. Tan PC, Khine PP, Vallikkannu N, Zawiah SZ. Promethazine compared with metoclopramide for hyperemesis gravidarum: a randomized controlled trial. **Obstet Gynecol.** 2010;115:975–981.
3. Levine MG, Esser D. Total parental nutrition for the treatment of severe hyperemesis gravidarum: maternal nutritional effects and fetal outcome. **Obstet Gynecol.** 1988;72:102–107.
4. Nelson-Piercy C. Treatment of nausea and vomiting in pregnancy: when should it be treated and what can be safely taken? **Drug Saf.** 1998;19:155–164.
5. Taylor R. Successful management of hyperemesis gravidarum using steroid therapy. **QJM.** 1996;89:103–107.
6. Jueckstock JK, Kaestner R, Mylonas I. Managing hyperemesis gravidarum: a multimodal challenge. **BMC Med.** 2010;8:46.
7. Simon EP, Schwartz J. Medical hypnosis for hyperemesis gravidarum. **Birth.** 1999;26:248–254.

Related policies

Australian College of Midwives. **National Midwifery Guidelines for Consultation and Referral.** 2013.3rd Edition.

Related WNHS policies, procedures and guidelines

KEMH. Hyperemesis Gravidarum: management of

Keywords:	Hyperemesis gravidarum, hyperemesis, KEMH		
Document owner:	Obstetrics, Gynaecology and Imaging Directorate		
Author / Reviewer:	Clinical Midwifery Manager CMP		
Date first issued:	October 2008		
Last reviewed:	September 2016	Next review date:	September 2019
Endorsed by:	OGCCU Management Committee	Date:	
Standards Applicable:	NSQHS Standards: 1  Clinical Care is Guided by Current Best Practice 4-  Medication Safety; 6-  Clinical Handover; 9  Clinical Deterioration,		
<p>Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.</p>			