



CLINICAL PRACTICE GUIDELINE
NEWBORN EMERGENCY TRANSPORT SERVICE (NETS WA)

Bronchiolitis

This document should be read in conjunction with the [Disclaimer](#)





Definition: viral infection of the lower respiratory tract manifesting as respiratory distress (and apnoea, especially in the young infant).

Management

- Provide oxygen as needed (Cot O₂ preferred to nasal prongs).
- CPAP if required. May be helpful in managing apnoea.
- Loading dose of IV Caffeine (20 mg/kg) if required.
- **Hypertonic Saline** (3%) has been shown to reduce airway oedema / increase mucous clearance and decrease secretion viscosity. Consider administering 2mLs 3% NaCl via nebuliser at referring hospital. Repeat as required.
- **Nebulised Adrenaline** has shown positive effects on short term measures including improved clinical score and reducing admission rates in the first 24 hours of care. Dose:
 - Using 1:1000 ampoule: 0.5 mL/kg/dose. Dilute to 2-4 mL with saline.
 - Using Respirator Solution: 0.05 mL/kg/dose.

Both Hypertonic Saline and Nebulised Adrenaline can only be administered at point of referral.

- Intubation and ventilation **after discussion with the on-call consultant**. Babies often deteriorate after intubation, and copious secretions may make ventilation difficult.
 - Sedation will be required.
 - Babies are prone to gas trapping.
 - Aim for lower rate, longer Ti (0.5-1.0) and Te, to allow better oxygenation and CO₂ removal but keep Ti < Te.
 - PEEP should generally be lower to avoid gas trapping, but higher PEEP may be needed in cases of atelectasis.
 - Regular ETT suction to prevent ETT occlusion.
- Nil by mouth, IV fluids at 2/3 maintenance.
- Start on antibiotics (Amoxicillin / Gentamicin).
- CXR (if available).

Document owner:	Neonatal Directorate Management Committee		
Author / Reviewer:	Neonatal Directorate Management Committee		
Date first issued:	August 2009		
Last reviewed:	1 st July 2017	Next review date:	1 st July 2020
Endorsed by:	Neonatal Directorate Management Committee	Date endorsed:	26 th September 2017
Standards Applicable:	NSQHS Standards: 1  Governance, 4  Medication Safety, 6  Clinical Handover, 9  Clinical Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.			