



CLINICAL PRACTICE GUIDELINE
NEWBORN EMERGENCY TRANSPORT SERVICE (NETS WA)

Fluid Management

This document should be read in conjunction with the [Disclaimer](#)

- Type and volume of fluid depends on infant’s gestational age and disease process.
- Typical glucose requirements for term neonates: 4-6 mg/kg/min and preterm neonates: 6-8 mg/kg/min.
- As a guide, use:
 - Normal saline for fluid expansion.
 - 10% glucose + 0.22% saline (> 34 weeks).
 - 7.5% glucose + 0.22% saline (27-34 weeks).
 - 5% glucose (≤ 27 weeks).
- Maintenance fluids for term neonates are usually approximately 60 mL/kg/day on Day 1.
 - Extreme preterm babies usually require much more (100-120 mL/kg/day) due to increased insensible fluid losses.
 - Asphyxiated infants are often fluid restricted to approximately 40mL/kg/day. In these circumstances, **the percentage concentration of glucose MUST be increased, as avoidance of hypoglycaemia is more important than fluid restriction.**

Document owner:	Neonatal Directorate Management Committee		
Author / Reviewer:	Neonatal Directorate Management Committee		
Date first issued:	August 2009		
Last reviewed:	1 st July 2017	Next review date:	1 st July 2020
Endorsed by:	Neonatal Directorate Management Committee	Date endorsed:	26 th September 2017
Standards Applicable:	NSQHS Standards: 1 Governance, 4 Medication Safety, 6 Clinical Handover, 9 Clinical Deterioration		

**Printed or personally saved electronic copies of this document are considered uncontrolled.
Access the current version from the WNHS website.**