



**CLINICAL PRACTICE GUIDELINE**

Guideline coverage includes NICU KEMH, NICU PCH and NETS WA

# Cardiac: Post-Operative Handover

This document should be read in conjunction with the [Disclaimer](#)

Most babies undergoing cardiac surgery will have their immediate post-op care in PCC with most returning to NICU as early as 24 hours later. This Guideline applies whether the baby returns from PCC or directly from Theatre.

## Prior to Arrival

Set up the bed space with appropriate monitoring equipment and drug infusions and ventilator settings as notified by theatre.

## Handover

Handover must follow the **iSoBAR** format.

The anaesthetist and NICU consultant and/or senior registrar, registrar and appropriate neonatal nurses should be present. All non-essential staff should move away and everyone should listen carefully and quietly to the handover.

<b>Identity</b>	Patient name and UMRN.
<b>Situation</b>	Describe the reason for handover.
<b>Observations</b>	TPR, BP, CVP, blood gas and haemoglobin. Ventilation settings and current infusions.
<b>Background</b>	Brief salient pre op status. <b>Surgery</b> <ul style="list-style-type: none"> <li>• Intra/post op echo details (if done).</li> <li>• Details of procedure.</li> <li>• Intra-operative surgical problems/complications.</li> </ul> <b>Anaesthesia</b> <ul style="list-style-type: none"> <li>• Itemise any ETT, vascular and surgical drain manipulations and difficulties.</li> <li>• Analgesia.</li> <li>• Blood losses and Fluid/blood product administration.</li> <li>• Any arrhythmia details.</li> </ul>
<b>Agree a plan</b>	Given the situation agree what needs to happen.
<b>Read back</b>	Confirm shared understanding.

## Following Handover

- Patient is transferred onto a NICU ventilator with immediate assessment of chest movement, air entry, end-tidal CO<sub>2</sub> and SaO<sub>2</sub>.
- Chest drains should be connected to suction (15-20cm H<sub>2</sub>O).

- Transfer transport monitoring to bedside monitoring. Invasive systemic BP and CVP monitoring is recommended.
- Review infusions concentrations. If changing inotrope infusion use the 'double pumped', i.e. the original infusion should only be stopped once the new infusion has 'hit'. You will be able to tell this when the BP rises.
- Medical and nursing staff should thoroughly examine the patient.
- ABG should be taken within 10-15 minutes of admission.
- FBC/ U+E/ Ca/ Mg/ coagulation profile should be checked.
- X-ray to check ETT, NGT, drain and line positions and lung and heart status.
- An ECG should be considered.

## Parents

Once surgeon has spoken to the parents and if patient is stable enough parents should be encouraged to see their child as soon as possible.

**Registrar or senior registrar should write all the above details in notes.**

Document owner:	Neonatal Directorate Management Committee		
Author / Reviewer:	Neonatal Directorate Management Committee		
Date first issued:	May 2013		
Last reviewed:	24 <sup>th</sup> November 2016	Next review date:	24 <sup>th</sup> November 2019
Endorsed by:	Neonatal Directorate Management Committee	Date endorsed:	16 <sup>th</sup> December 2016
Standards Applicable:	NSQHS Standards: 1  Governance, 5  Patient ID/Procedure Matching, 6  Clinical Handover, 9  Clinical Deterioration		
<b>Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.</b>			