



CLINICAL GUIDELINE

**Antenatal Renal and Urological Anomalies**

|                       |                           |
|-----------------------|---------------------------|
| <b>Scope (Staff):</b> | Nursing and Medical Staff |
| <b>Scope (Area):</b>  | KEMH Postnatal Wards      |

This document should be read in conjunction with this [DISCLAIMER](#)

## Background

Abnormalities of the foetal genito-urinary system are identified by ante-natal ultrasonography in 0.5% of pregnancies<sup>1</sup>. Antenatal Renal Pelvis Dilatation (RPD) affects approximately 1% of fetuses<sup>2</sup>. Elevated RPD indicating the presence of hydronephrosis is the most common urinary tract finding on routine ante-natal scanning and clinical sequelae may range from no functional disturbance, through to acute and chronic renal failure, depending on the underlying cause. Whilst many instances of antenatally diagnosed hydronephrosis are associated with a normal renal prognosis, underlying abnormalities such as spina bifida, posterior urethral valves, obstruction at the pelvi-ureteric junction, duplex system and other congenital anomalies may have significant implications with respect to renal functional outcomes. Foetal RPD measures of > 15 mm have demonstrated an association with renal pathology such as urinary tract obstruction<sup>2</sup>. This diagnosis implies a need for early surgical intervention in most cases. The presence of unexpected renal fluid collections on ultrasound (e.g. urinoma, renal cysts, distended bladder) should prompt consideration of a co-existent renal tract anomaly<sup>3</sup>.

Defining the extent and severity of hydronephrosis identified on antenatal ultrasound has been confounded by physiological variation in growth occurring normally during gestation, and consequent to differences in definition and diagnostic criteria. The most commonly used method for diagnosing prenatal hydronephrosis is the assessment of the antero-posterior diameter of the renal pelvis<sup>2</sup>. Unfortunately, opinion varies regarding which degrees of renal dilatation require investigation with ultrasound or other modalities in the post-natal period.

The use of antibiotic prophylaxis, surgical intervention for (e.g.) vesico-ureteric reflux and long-term monitoring strategies are also the focus of debate. Meta-analysis of outcomes for mild-moderate RPD have demonstrated an increasing risk of obstruction / VUR with increasing mean foetal pelvis diameter increased from 5mm to 15mm<sup>2</sup>. Other meta-analyses have indicated that milder degrees of pelvis dilatation were more likely to stabilise or improve<sup>4</sup>, while moderate to severe cases present a significant risk of association with an underlying pathology<sup>5</sup>.

## Management of Infants with Mild Antenatal RPD

Infants with a renal pelvis diameter of **5-10mm** on ultrasound are considered to have mild renal dilatation. Infants in whom mild renal dilatation is identified and is persistent on antenatal scan should be managed as follows:

- Refer to Nephrology Unit using [referral form](#).
- Commence Bactrim prophylaxis, 1mL orally daily. To continue until renal US and review.

- Renal Unit Secretary will arrange ultrasound and outpatient appointment and this will be sent to parents. These may not necessarily occur on the same day.

### Management of Infants with Moderate-Severe Antenatal RPD

Infants with a renal pelvis diameter of **> 10mm** on ultrasound are considered to have moderate to severe (> 15mm) renal pelvis dilatation. Infants in whom moderate or severe dilatation is present should be managed as follows:

- Ring the on-call Consultant Nephrologist at PCH via Switchboard.
- Commence Bactrim prophylaxis, 1mL orally daily. To continue until renal US and review.
- Complete renal [follow-up referral](#) template and/or arrange for other investigations as requested by the Nephrologist. Urgent renal ultrasounds can generally be obtained at short notice at KEMH if discussed with the radiologist and ultrasonographer on call **after** discussion with the PCH nephrologist or PNW paediatric consultant.

### Management of Infants with Other Renal Anomalies


Infants, in whom other renal tract anomalies have been identified in utero, should be assessed clinically and discussed with the paediatric consultant on call for post-natal wards prior to consulting Renal or other specialties.

Antibiotic Prophylaxis for renal tract dilatation or anomalies

Antibiotic prophylaxis should be initiated for all infants with renal pelvis dilatation, pending outpatient ultrasound and clinical review by PCH Nephrology staff. **Bactrim (co-trimoxazole) 1mL orally daily** should be used unless specifically ordered otherwise by the paediatric / renal consultant. The medication should be continued after discharge until renal review, at which time a decision will be made regarding its continuation.

### Urology and Renal Referral Form

This document can be made available in alternative formats on request for a person with a disability.

|   |  |                   |                             |
|---|--|-------------------|-----------------------------|
| File Path:  |  |                   |                             |
| Document Owner:   | Neonatology  |                   |                             |
| Reviewer / Team:  | Neonatal Coordinating Group  |                   |                             |
| Date First Issued:  | August 2011  | Last Reviewed:    | 24 <sup>th</sup> March 2020 |
| Amendment Dates:  |  | Next Review Date: | 24 <sup>th</sup> March 2023 |
| Approved by:  | Neonatal Coordinating Group  | Date:             | 28 <sup>th</sup> April 2020 |
| Endorsed by:  | Neonatal Coordinating Group  | Date:             |                             |
| Standards Applicable:   | NSQHS Standards:  |                   |                             |
| <b>Printed or personally saved electronic copies of this document are considered uncontrolled</b> |  |                   |                             |