






ADULT

IMIQUIMOD

This document should be read in conjunction with this [DISCLAIMER](#)

RESTRICTED Medication

Presentation	Cream sachet: 5% (50mg/g)
Dose	<u>Condylomata acuminata (Genital warts)</u> Topical: 3 nights per week (For up to 16 weeks)
Administration	Patient to self-apply before bedtime. Before applying, wash affected area with mild soap and water and allow to dry. Wash off with mild soap and water about 6-10 hours after application. Wash hands before and after use; avoid bathing/showering after application.
Pregnancy	1st Trimester: Consider alternative 2nd Trimester: Consider alternative 3rd Trimester: Consider alternative
Breastfeeding	Considered safe to use
Clinical guidelines and policies	WNHS Guideline: Sexually Transmitted Infections (STI)
References	Australian Medicines Handbook. Imiquimod. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2017 [cited 2017 Sept 20]. Available from: https://amhonline.amh.net.au/ The Royal Women's Hospital. Imiquimod. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2016 [cited 2017 Sept 20]. Available from: https://thewomenspbmg.org.au/

Keywords:	Imiquimod, Aldara, Genital warts, Sexually transmitted infection		
Publishing:	<input checked="" type="checkbox"/> Intranet <input checked="" type="checkbox"/> Internet		
Document owner:	Chief Pharmacist		
Author / Reviewer:	Pharmacy Department		
Date first issued:	September 2017	Version:	3.0
Last reviewed:	September 2017	Next review date:	Sep 2020
Endorsed by:	Medicines and Therapeutics Committee	Date:	Dec 2017
Standards Applicable:	NSQHS Standards: 1  Clinical Care is Guided by Current Best Practice 3  Infection Control 4  Medication Safety		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the WNHS website.			

© Department of Health Western Australia 2017