



ADULT MEDICATION GUIDELINE

PARECOXIB

Scope (Staff): All WNHS Staff

Scope (Area): Obstetrics and Gynaecology

This document should be read in conjunction with the [Disclaimer](#).

Quick Links

[Dose](#)

[Administration](#)

[Monitoring](#)

[Pregnancy and
Breastfeeding](#)

Restrictions

[Formulary: Unrestricted](#)

Medication Class

Nonsteroidal Anti-inflammatory Drug (NSAID)

Presentation

Vial: 40mg

Storage

Store at room temperature, below 25°C

Dose

Schedule 8 Medication

Postoperative Pain

IV/IM:

40mg as a single dose

In patients >65yrs and <50kg or with moderate hepatic impairment:

20mg as a single dose

Administration

Step 1 Reconstitution:

IV/IM

Reconstitute the vial with 2 mL of sodium chloride 0.9%, glucose 5% or glucose 5% in sodium chloride 0.45%; the solution is clear and colourless

Step 2 Administration:

IV injection

Inject over a few seconds

IM injection

Inject slowly and deep into the muscle

Monitoring

Renal: Avoid use if CrCl <25 mL/minute

Hepatic: Contraindicated in severe impairment; reduce dose in moderate impairment (Child-Pugh score 7–9) as elimination is decreased

Elderly: Reduce dose for elderly women <50 kg

Pregnancy

1st Trimester: Consider alternative

2nd Trimester: Consider alternative

3rd Trimester: Consider alternative

For more information, please contact [KEMH Obstetric Medicines Information Service](#).

Breastfeeding

Considered safe to use

Comments

Regular oral NSAIDs may be commenced as soon as 1 hour after IV parecoxib or 2 hours after rectal diclofenac.

Related Policies, Procedures & Guidelines

WNHS Clinical Practice Guidelines:

[The Hitchhikers Guide to Obstetric Anaesthesia](#)

WNHS Pharmaceutical and Medicines Management Guidelines:

[WNHS Medication Administration Guideline](#)

[WNHS Medication Management Guideline "Return and Disposal of Medications"](#)

References









Australian Medicines Handbook. Parecoxib. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2021 [cited 2021 July 20]. Available from: <https://amhonline.amh.net.au/>

The Royal Women's Hospital. Parecoxib. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2021 [cited 2021 July 20]. Available from: <https://thewomenspbmg.org.au/>

Society of Hospital Pharmacists of Australia. Parecoxib. In: Australian Injectable Drugs Handbook [Internet]. [St Leonards, New South Wales]: Health Communication Network; 2021 [cited 2021 July 20]. Available from: <http://aidh.hcn.com.au>

MIMS Australia. Parecoxib (Dynastat®). In: MIMS Online [Internet]. St Leonards (New South Wales): MIMS Australia; 2021 [cited 2021 July 20]. Available from: <https://www.mimsonline.com.au>

MIMS Australia. Voltaren® suppositories. In: MIMS Online [Internet]. St Leonards (New South Wales): MIMS Australia; 2021 [cited 2021 July 20]. Available from: <https://www.mimsonline.com.au>

Keywords					
Document Owner:	Chief Pharmacist				
Author/ Reviewer	KEMH Pharmacy Department				
Version Info:	4.0				
Date First Issued:	20/09/2007	Last Reviewed:	07/09/2021	Review Date:	07/09/2024
Endorsed by:	Medicines and Therapeutics Committee			Date:	16/09/2021
NSQHS Standards Applicable:	<input checked="" type="checkbox"/>  Std 1: Clinical Governance <input type="checkbox"/>  Std 2: Partnering with Consumers <input type="checkbox"/>  Std 3: Preventing and Controlling Healthcare Associated Infection <input checked="" type="checkbox"/>  Std 4: Medication Safety		<input type="checkbox"/>  Std 5: Comprehensive Care <input type="checkbox"/>  Std 6: Communicating for Safety <input type="checkbox"/>  Std 7: Blood Management <input type="checkbox"/>  Std 8: Recognising and Responding to Acute Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from WNHS HealthPoint.					

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability.

© Women and Newborn Health Service 2021

Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.