

Ultrasound patient information

Saline Infusion Sonography (SIS) and HyCoSy care

Date of Procedure:	

It is normal to have some cramping or period-like pains following your procedure. Use of local warmth (such as a hot-water bottle or wheat bag) is usually sufficient to help.

You may take analgesics such as paracetamol, Nurofen, Naprogesic, Ponstan or Advil, if necessary for the pain.

You will probably notice a dark discharge from the vagina, due to the Betadine antiseptic used. This usually stops after a day or two.

You may experience a small amount of vaginal spotting or bleeding in the day or two following. This usually settles down without treatment.

No sexual intercourse for the 24 hours following the procedure.

If you become unwell with abdominal pains, unusual or smelly discharge from the vagina, fevers and chills or loss of appetite, then you must contact your Doctor or come to the Emergency Centre at King Edward Memorial Hospital (telephone number 6458 1431).

Your next step in management planning will be made by your referring doctor with you at your next review appointment.

For further information about ultrasounds at KEMH

Ultrasound Department - (08) 6458 2830



This document can be made available in alternative formats on request.

© North Metropolitan Health Service - Aug 2018 Produced by:

Women and Newborn Health Service King Edward Memorial Hospital 374 Bagot Road, Subiaco WA 6008 Telephone: (08) 6458 2222 www.kemh.health.wa.gov.au

Disclaimer: The advice and information contained herein is provided in good faith as a public service. However the accuracy of any statements made is not guaranteed and it is the responsibility of readers to make their own enquiries as to the accuracy, currency and appropriateness of any information or advice provided. Liability for any act or omission occurring in reliance on this document or for any loss, damage or injury occurring as a consequence of such act or omission is expressly disclaimed.